

FACTSHEET

Veterinary Services

United States
Department of
Agriculture

Animal and
Plant Health
Inspection
Service

December 1998

Screwworm

Screwworm is the common name of a pest native to the tropical areas of North, South, and Central America that causes extensive damage to domestic livestock and other warmblooded animals. The larvae of these pests feed on the raw flesh of the host animal. Rare human cases have been reported.

The United States has been free of screwworm since 1966. The U.S. livestock industry could suffer \$750 million in production losses annually if this pest were reintroduced to the United States.

The Pest and Its Life Cycle

The screwworm is an insect that, in its adult stage, is about twice the size of the common housefly. It has orange eyes and a blue-grey or grey body with three dark stripes running down its back.

After mating, the female screwworm fly lays her eggs in the open wounds of animals. One female can lay up to 400 eggs at a time and as many as 2,800 eggs during its 31-day lifespan. These eggs can hatch into larvae in as little as 12 hours.

Screwworm larvae grow by feeding on the flesh of living animals and can grow to be over one-half inch within 5 to 7 days after hatching. The full-grown larvae then drop from the wound and tunnel into the soil, where they form protective cases to house themselves while they pupate. Adult screwworm flies emerge from these pupal cases and are ready to mate again within 3 to 5 days.

Clinical Signs of Screwworm Infestation

Animals are infested when the eggs hatch in the wound of an animal and the larvae feed on the animal's flesh. Wounds that may become infested by screwworm include those caused by feeding ticks, bites of vampire bats, castration, dehorning, branding, shearing, wire cuts, sore mouth in sheep, and shedding of the velvet in deer. Navels of newborn mammals are a common site for screwworm infestation. It is very difficult to see early stages of screwworm larvae feeding in a wound; only slight

movement may be observed. As the larvae feed, the wound is gradually enlarged, becoming wider and deeper.

Usually at this stage, additional screwworm flies have deposited eggs, resulting in a multiple infestation. A bloody discharge often exudes from the infested wounds, and a distinct odor may be detected. In some cases, the openings in the skin may be small with extensive pockets of screwworm larvae beneath.

Animals with screwworm infestations may die in 7 to 14 days if wounds are not treated to kill the larvae, especially in cases of multiple infestations. As many as 3,000 larvae may be found in a single wound. Death results from toxicity and/or secondary infection.

Infested animals usually exhibit discomfort, may go off feed, and produce less milk. Typically, these animals also will separate themselves from the rest of the flock or herd and seek shady or secluded areas to lie down.

Control

A screwworm infestation is treated with topical application of approved chemicals to kill the larvae. Wounds should be treated for 2 to 3 consecutive days to ensure that all larvae have been killed. The larvae should be removed from the wounds using tweezers.

Treating wounds and spraying or dipping animals with an approved organophosphate insecticide will provide protection against screwworm for 7 to 10 days. However, the most effective way to control screwworm infestation is through eradication.

History of the U.S. Screwworm Eradication Program

As early as 1825, U.S. Western States were reporting serious problems with screwworm. By the 1930's, screwworm had spread to the Southeastern States, where livestock producers were losing \$400 million annually.

In the early 1950's, the U.S. Department of Agriculture's (USDA) Agricultural Research Service developed a new method to help eradicate the pest using a form of biological control. During the pupal

stage of the fly's life cycle, the pupae are subject to gamma radiation designed to leave the fly sexually sterile. The flies are artificially raised in a production plant (originally located in Mission, TX) and then released by aircraft over infested areas. When the sterile males mate with females of the native fly populations, no offspring result. With fewer fertile mates available in each succeeding generation, the fly, in essence, breeds itself out of existence.

This sterile insect technique was successfully tested in a field trial on the Dutch island of Curaçao in 1954 and then used operationally in Florida in 1957. By 1959, screwworm had been eradicated from the entire Southeastern United States. The technique was next applied to the more extensively infested Southwest in 1962. Self-sustaining screwworm populations were eliminated from the United States by 1966. A barrier zone of sterile flies was established along the 2,000-mile-long U.S.–Mexico border to prevent reinfestation from Mexico.

However, constant reinfestation from migrating flies or larvae carried by animals, which are then transported by people, remained a problem. Therefore, in 1972, the United States–Mexico Joint Commission for the Eradication of Screwworm was formed with the goal of eliminating the pest from Mexico and pushing the sterile fly barrier to the Isthmus of Tehuantepec, just north of Guatemala. A new sterile screwworm plant at Tuxtla Gutierrez, Chiapas, Mexico, was dedicated in 1976, replacing the former production plant in Mission, TX, which closed in January 1981. As a result of the Commission's efforts, Mexico was officially declared free of the pest in 1991.

APHIS has also been cooperating with Central America to eradicate screwworm from those countries and ultimately to establish and maintain a permanent sterile fly barrier at the Darien Gap between Panama and Colombia. The Darien Gap is the narrowest geographical region in southern Panama, stretching only 102 miles. The region is mainly jungle and has no roads, so there is minimal risk of infested animals being transported by land from South America into Panama.

In addition, as part of the program's overall strategy, a new sterile-fly rearing facility in Panama, to be established around 2003, will replace the existing one in Mexico. Locating the new facility in

Panama, an area where screwworm has not been eradicated, will reduce the risk of reinfestation of the United States through accidental release of fertile flies.

To date, Belize, Guatemala, El Salvador, and Honduras have been declared free of screwworm. Currently, screwworm program officials are focusing their efforts on eradicating the pest from Nicaragua and Costa Rica. Eradication activities include regulation of cattle movement, treatment of wounds, and release of sterile flies.

Threat to U.S. Cattle Producers

Although the screwworm program has been very successful, limited outbreaks have occurred. In 1992 and 1993, the program's resources were strained by introduced outbreaks in Mexico. Both outbreaks were contained and eradicated, but at a cost of several million dollars. In that outbreak, the closest case identified was only 121 miles from the U.S. border.

In 1997, a screwworm larva was detected on a dog shipped to San Antonio, TX, from a military base in Panama. Fortunately, a quick-thinking private veterinary practitioner detected and collected the larva. State and Federal animal health authorities immediately took appropriate measures to track the dog's movement into the United States, disinfect sites, and prevent the spread of the pest.

Report Suspicious Cases

These incidents remind us that screwworm is still a real threat to U.S. livestock. Veterinarians and livestock owners who suspect an animal may be infested with screwworm larvae should immediately contact State or Federal animal health authorities.

For more information, contact
USDA, APHIS, Veterinary Services
Emergency Programs
4700 River Road, Unit 41
Riverdale, MD 20737–1231
Telephone: (301) 734–8073
Fax: (301) 734–7817

Current information on animal diseases and suspected outbreaks is also available on the Internet. Point your Web browser to <http://www.aphis.usda.gov> to reach the APHIS home page.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720–2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326–W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250–9410 or call (202) 720–5964 (voice and TDD). USDA is an equal opportunity provider and employer.